

UL'YANOV, N.A., kand.tekhn.nauk

Evaluating tractive properties of wheel drives in earthmoving machines.  
Stroi.i dor.mashinostr. 5 no.3:16-20 Mr '60. (MIRA 13;6)  
(Traction engines)  
(Earthmoving machinery)

MIKHAYLOV, B.I., inzh.; UL'YANOV, N.A., kand.tekhn.nauk

Automatic adjustment of motor grader operations. Stroi.i dor.  
mashinostr. 5 no.7:6-7 JI '60. (MIRA 13:7)  
(Automatic control)  
(Graders (Earthmoving machinery))

UL'YANOV, N. A., dotsent, kand. tekhn. nauk

Choice of parameters and operating conditions of a wheel-mounted motor of continuous earthmovers with cutting blades.  
Sbor. trud. MISI no.39:268-274 '61. (MIRA 16:4)

(Earthmoving machinery)

UL'YANOV, Nikolay Aleksandrovich, kand. tekhn. nauk; BAZANOV, A.F.,  
kand. tekhn. nauk, retsenzent; KONONENKO, M.A., inzh., red  
SAVEL'YEV, Ye.Ya., red. izd-va; SMIRNOVA, G.V., tekhn. red.

[Fundamentals of the theory and design of wheeled tractors  
for excavating machinery] Osnovy teorii i rascheta kolesnogo  
dvizhitelia zemleroiinykh mashin. Moskva, Mashgiz, 1962.  
206 p. (MIRA 16:4)

(Tractors--Design and construction)  
(Excavating machinery)

UL'YANOV, N.A., kand.tekhn.nauk

Method of making traction computations for rollers on  
pneumatic tires. Stroi. i dor. mash. 7 no.8:15-16 Ag '62.  
(MIRA 15:9)

(Rollers (Earthwork))

ALEKSEYEVA, T.V., kand. tekhn. nauk; ARTEM'YEV, K.A., kand. tekhn. nauk; BROMBERG, A.A., prof.; VOYTSEKHOVSKIY, R.I., inzh.; UL'YANOV, N.A., kand. tekhn. nauk; Primal uchastiye KONONENKO, M.A., inzh.; FEDOROV, D.I., kand. tekhn. nauk, retsenzent.

[Machines for earthwork; theory and calculation] Mashiny dlia zemlianykh rabot; teoriia i raschet. [By] T.V. Alekseeva i dr. Izd.2., perer. i dop. Moskva, Izd-vo "Mashinostroenie," 1964. 467 p. (MIRA 17:5)

UL'YANOV, N.G.

Testing an experimental hydraulic clutch in a ZIS-150 car. Sborn.trud.  
lab.preb.bystr.mash. 3:205-213 '53. (MIRA 9:9)  
(Automobile--Transmission devices)

VASIL'YEVA, N.N.; UL'YANOV, N.K.

Geobotanical studies as a method of prospecting for ore deposits  
in central Kazakhstan. Inform.sbor.VSEGEI no.50:83-94 '61.  
(MIRA 15:8)

(Kazakhstan--Prospecting) (Kazakhstan--Phytogeography)



TSYKUNKOVA, N.A.; UL'YANOV, N.K.

Occurrences of metals in eluvial and talus formations of some ore  
deposits in central Kazakhstan. Inform.sbor.VSEGEI no.50:71-81  
'61. (MIRA 15:8)

(Kazakhstan—Metals, Rare and minor)  
(Kazakhstan—Nonferrous metals)

MAROCHKIN, N.I., glav. red.; MARKOVSKIY, A.P., zam. glav. red.;  
UL'YANOV, N.K., zam. glav. red.; GANESHIN, G.S., red.;  
ZAYTSEV, I.K., red.; KNIPOVICH, Yu.N., red.; KULIKOV, M.V., red.;  
LABAZIN, G.S., red.; LUR'YE, M.L., red.; SIMONENKO, T.N., red.;  
SPIZHARSKIY, T.N., red.; STERLIN, D.Ya., red.; TATARINOV, P.M., red.;  
BELYAKOVA, Ye.Ye., nauchnyy red.; MAKRUSHIN, V.A., tekhn. red.

[Yearbook of the results of studies by the All-Union Geological  
Institut] Ezhegodnik po rezul'tatam rabot VSEGEI. Leningrad,  
Otdel nauchn.-tekhn. informatsii, 1961. 203 p. (Leningrad.  
Vsesoiyuznyi geologicheskii institut. Informatsionnyi sbornik,  
no.49.) (MIRA 15:6)

(Geology)

UL'YANOV, N.N., inzh.; SHPORKHUN, V.I., inzh.

Distributing device for the refluxing of packed columns. Khim.  
mashinostr. no.3:3-4 My-Je '63. (MIRA 16:11)

ARUTYUNYAN, B.Sh.; BORISOV, V.M.; ZHEPLINSKIY, B.M.; MESROPYAN, N.N.;  
MESHCHERYAKOV, N.F.; UL'YANOV, N.S.

Apparatus for the destruction of flotation froth. Khim. prom.  
no.2:146-147 F '63. (MIRA 16:7)

(Flotation)

10

CONCENTRATION OF PHOSPHORITES. N. S. Ulanov. *Bull. acad. sci. U. R. S. S., Class sci. math. nat., Ser. chim.* 1938, No. 1, 60-73 (in English 74).—The primary method of concn. now in practice yielded, at best, a concentrate contg. 25-65% of  $P_2O_5$  and 6-7% of  $R_2O_3$ . An exptl. flotation of preliminary calcined phosphorite yielded a concentrate contg.  $P_2O_5$  29.2 and  $R_2O_3$  4.8% (the same phosphorite ore as above) and in some instances contg. even 30-1% of  $P_2O_5$  and 5.5% of  $R_2O_3$  (different ore). Of all concn. methods, flotation yielded the best results. Six references.

A. A. Podzornov

ASM-55A METALLURGICAL LITERATURE CLASSIFICATION

100000 00 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 00

1ST AND 2ND ORDERS																										3RD AND 4TH ORDERS																									
PROCESSING AND PROPERTIES INDEX																																																			
18																																																			
<p>Recent work and present problems in enriching phosphorite. N. S. Ul'yanov. J. Chem. Ind. (U. S. S. R.) ; 1960 No. 11, 23-25 (1961). See C. A. 32, 8700d. H. M. L.</p>																																																			
<p>ASM-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>																																																			
<p>1ST AND 2ND ORDERS</p>																																																			
<p>3RD AND 4TH ORDERS</p>																																																			

1ST AND 2ND CODES										PROCESS AND PROPERTIES INDEX										3RD AND 4TH CODES										
																														<b>18</b>
<p><i>Cd</i></p> <p>Flotation on a semiproduction scale, with preliminary roasting of the Ryazan-Akvilov ore from the Egor'ev ore deposits. N. S. Ul'yangya, V. M. Vidonov and N. V. Maharenko. <i>Khimiya Nefti i Uglekhimii</i>, Gubkentizn, Nernykh Rd., Shornik Rabot. Nauch. Inst. Udobreniyam, Izvestiya Akad. Nauk SSSR, No. 150, 50-51, 1963, 1 p. — Roasting the 73; Khim. Referat. Zhur. 1940, No. 6, 84. — Roasting the washed Ryazan-Akvilov ore, before flotation, in a Poly-washed furnace for 40-50 min. improves considerably the quality of the flotation concentrate, bringing the content of P<sub>2</sub>O<sub>5</sub> to 30-1% and K<sub>2</sub>O to 4.3-6.5%. Duration of the basic flotation is 16 min. and purification of the concentrate 8 min. The amts. of the reagents used are soda 6.0-0.5, acidole 5, kerosene 5 and water glass 0.75 kg./ton.</p> <p align="right">W R. Hean</p>																														
<p align="center">ABX-SLA METALLURGICAL LITERATURE CLASSIFICATION</p> <p align="right">CLASSIFICATION</p>																														
<p>CLASSIFICATION</p>																														

15

Flotation of phosphorite ore of the Portland horizon of the Egor'ev ore deposits. N. S. Ulyanov and V. M. Kuznetsovskaya. *Obogatitel'noye Priblizheniye* i *Sovetskaya Rud. Shkola* Nauch. Inst. Uchebnik i *Iskustvennyy* Ya. V. Samoilov 1939, No. 15, 93-103; *Khim. Refort. Zhur.* 1940, No. 6, 84-8. — Washing the Portland ore on a 0.25-mm. screen produced a +0.25 mm. concentrate contg.  $P_2O_5$  31.9,  $R_2O_3$  9.4 and insol. residue 17.94%. The optimum fineness of the ore is 100 mesh, with not more than 7% of the residue remaining on the screen. The flotation reagents were: carboxylic acids 1.7, water glass 1.0 and lime 1.0 kg./ton. In the pulp the ratio solid liquid was 1:4. After flotation, the concentrate contained  $P_2O_5$  27.5-27.9,  $R_2O_3$  3.9-4.5 and insol. residue 0.0-0.5%. The extn. was 91.1%. A 0.25-mm. flotation product produced a concentrate contg. 14-16% of  $P_2O_5$ , which can be used to produce phosphorite meal. W. R. Henn

ASB-51A METALLURGICAL LITERATURE CLASSIFICATION

FROM 1771814VW

12345678910111213141516171819202122232425262728293031323334353637383940414243444546474849505152535455565758596061626364656667686970717273747576777879808182838485868788899091929394959697989900

10111213141516171819202122232425262728293031323334353637383940414243444546474849505152535455565758596061626364656667686970717273747576777879808182838485868788899091929394959697989900



COMMON ELEMENTS										PROCESSING AND PREPARATION INDEX										1ST AND 2ND ORDER										3RD AND 4TH ORDER									
<p>4</p> <p>Flotation of the Upper Kama phosphorites ore complex. N. S. Ulyanov. <i>Obozrazheniye Fosforitov, Glandonilov i Svezhsk Rud, Shornik Rabot Nauch. Inst. Udobreniy i Ischislofungisidam Ya. V. Samolovs</i> 1939, No. 150, 103-20; <i>Khim. Referat. Zhur.</i> 1940, No. 6, 85. — Proper conditions for flotation were developed from lab. and semi-production expts. and a qual. and a quant. scheme for enriching the Upper Kama phosphorites preppt. The ore is ground to 150 mesh (with a 8-10% residue in the sieve). Flotation can be carried out without settling of the fine slimes, with a 3-fold purification of the concentrate and a single purification of the tailings. After the 3rd purification the concentrate contained <math>P_2O_5</math> 27.5-27.8 and <math>R_2O_3</math> 4.37-4.70%. The <math>P_2O_5</math> extn. was 93.4-95.8% of the washed concentrate. The main flotation process took 13.5 min., purification of the concentrate 13.0-18.5 min. and purification of the tailings 13 min. The reagents used for the main flotation were carboxylic acid 2.00-3.00 and carboxylic acid salt 0.20 kg./ton. Purification of the tailings required carboxylic acid 1.54-1.56 and carboxylic acid salt 0.14 kg./ton. No water glass was required.</p>										<p>18</p>										<p>The Fahrwald machines are suitable for flotation of the Upper Kama phosphorites. W. R. Henn</p>																			
ASH-51A METALLURGICAL LITERATURE CLASSIFICATION										FROM ROMANOV										FROM ROMANOV																			
147082 24										147082 24										147082 24																			
147082 24										147082 24										147082 24																			

1ST AND 2ND CODES																										3RD AND 4TH CODES																									
PROCESSES AND PROPERTIES INDEX																																																			
<p><i>Flotation of the Upper Kama phosphorite ores with a preliminary roasting. N. S. Ulyanov, V. M. Vidonov, and V. G. Konstantinov. <i>Oblastskhaya Fosforitovaya Gidrometallurgicheskaya V. V. Samoilova 1939, No. 180, 120-122; Khim. Referat. Zhur. 1940, No. 6, 85.</i> Washed concentrate, ground to 16-mm. mesh, was used for the preliminary roasting under semicon. conditions. The compn. of the concentrate was <math>P_2O_5</math> 22.3-23.79 and <math>R_2O_3</math> 8.46-9.31%. The roasting was done in Polysius furnaces having an output of 100-40 kg./hr. The losses were 7-8%. By flotation in a 12-chamber Fahrenwald machine 2 fractions were obtained, one of which (48% of the total) contained 30% <math>P_2O_5</math> and 4% <math>R_2O_3</math>, and the other (52% of the total) contained 21.1% <math>P_2O_5</math>. The filtration rate in a disk vacuum filter was 420 kg./sq. m., as compared with 56 kg./sq. m. with the unroasted ore. The reagents used were soda 6.0-6.5, acido 2, crude oil 0.6, kerosene 8.6 and water glass 0.2 kg./ton for the 1st purification and 0.15 kg./ton for the 2nd purification. The duration of the main flotation was 9 min., of the 1st purification 5.0-5.5 min., of the 2nd purification 8.0-8.5 min. and of the 3rd purification 10.0-16.5 min. The ratio solid:liquid in the pulp was 1:4. Cf. C. A. 36, 4679, and preceding abstr. W. R. Henn</i></p>																																																			
<p>ASB-5LA METALLURGICAL LITERATURE CLASSIFICATION</p>																																																			
<p>12000 27000 31000 32000 33000 34000 35000 36000 37000 38000 39000 40000 41000 42000 43000 44000 45000 46000 47000 48000 49000 50000 51000 52000 53000 54000 55000 56000 57000 58000 59000 60000 61000 62000 63000 64000 65000 66000 67000 68000 69000 70000 71000 72000 73000 74000 75000 76000 77000 78000 79000 80000 81000 82000 83000 84000 85000 86000 87000 88000 89000 90000 91000 92000 93000 94000 95000 96000 97000 98000 99000</p>																																																			

*Ch*

*18*

Flotation of phosphorite ore from the Shchigrov deposits. N. S. Ul'yanov and V. M. Rauchevskaya. (*Ogashchenie Polymetallov, Gidrometallurgiya i Seruykh Rud, Stroyak Nauch. Inst. Uchebniy i Issledovaniyskiy Ye. V. Samoilov* 1939, No. 150, 145-51; *Khim. Refrat. Zhur.* 1940, No. 6, 85-9.—The av. P<sub>2</sub>O<sub>5</sub> content in the Shchigrov deposits is 12%. The raw material is ground to 150-200 mesh. Carboxylic acids are used as reagents. A concentrate was obtained contg. P<sub>2</sub>O<sub>5</sub> 26.8-28.0 and R<sub>2</sub>O<sub>3</sub> 3.7-4.1%; tailings contained P<sub>2</sub>O<sub>5</sub> 3.57%; estn. was 75-80%.

W. R. Hene

18

*Ch*

Extraction of glauconite and phosphorite from the tailings of concentrating plants of the Egor'ev and Upper Kama deposits. N. S. Il'yangov. *Obezashchenie Fosforitov, Glaukonitov i Sernykh Rud, Shornik Rabot Nauch. Inst. L'obremiyam i Issledovaniyam Ya. V. Samoilova 1939, No. 150, 152-50; Khim. Referat. Zhur. 1940, No. 6, 86.*

Enriching phosphorite ore from the Egor'ev and Upper Kama deposits gives low extn. of  $P_2O_5$ . The tailings contain not only phosphorite, but also glauconite (hydrous aluminosilicate of Fe, K, Ca, Mg), which is used for softening hard water. The method used to obtain the glauconite concentrate from tailings includes condensation, classification, drying, bolting, magnoetic sepn. and roasting. The initial material, when ground to 60-5 mesh, contains a max. amt. of glauconite. The +0.6 mm. fraction, obtained during bolting, before the magnetic sepn., and the tailings of the sepd. fraction are used to enrich phosphorites. Flotation with dark naphtha soap gives a concentrate contg. 18-20%  $P_2O_5$ . Such enrichment increases the  $P_2O_5$  extn. from 65-70 to 85%. A method for enriching the Egor'ev deposits ore is described. W. R. Heun

ASAC-ELA METALLURGICAL LITERATURE CLASSIFICATION

1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2722 2723 2724 2725 2726 2727 2728 2729 2730 2731 2732 2733 2734 2735 2736 2737 2738 2739 2740 2741 2742 2743 2744 2745 2746 2747 2748 2749 2750 2751 2752 2753 2754 2755 2756 2757 2758 2759 2760 2761 2762 2763 2764 2765 2766 2767 2768 2769 2770 2771 2772 2773 2774 2775 2776 2777 2778 2779 2780 2781 2782 2783 2784 2785 2786 2787 2788 2789 2790 2791 2792 2793 2794 2795 2796 2797 2798 2799 2800 2801 2802 2803 2804 2805 2806 2807 2808 2809 2810 2811 2812 2813 2814 2815 2816 2817 2818 2819 2820 2821 2822 2823 2824 2825 2826 2827 2828 2829 2830 2831 2832 2833 2834 2835 2836 2837 2838 2839 2840 2841 2842 2843 2844 2845 2846 2847 2848 2849 2850 2851 2852 2853 2854 2855 2856 2857 2858 2859 2860 2861 2862 2863 2864 2865 2866 2867 2868 2869 2870 2871 2872 2873 2874 2875 2876 2877 2878 2879 2880 2881 2882 2883 2884 2885 2886 2887 2888 2889 2890 2891 2892 2893 2894 2895 2896 2897 2898 2899 2900 2901 2902 2903 2904 2905 2906 2907 2908 2909 2910 2911 2912 2913 2914 2915 2916 2917 2918 2919 2920 2921 2922 2923 2924 2925 2926 2927 2928 2929 2930 2931 2932 2933 2934 2935 2936 2937 2938 2939 2940 2941 2942 2943 2944 2945 2946 2947 2948 2949 2950 2951 2952 2953 2954 2955 2956 2957 2958 2959 2960 2961 2962 2963 2964 2965 2966 2967 2968 2969 2970 2971 2972 2973 2974 2975 2976 2977 2978 2979 2980 2981 2982 2983 2984 2985 2986 2987 2988 2989 2990 2991 2992 2993 2994 2995 2996 2997 2998 2999 3000 3001 3002 3003 3004 3005 3006 3007 3008 3009 3010 3011 3012 3013 3014 3015 3016 3017 3018 3019 3020 3021 3022 3023 3024 3025 3026 3027 3028 3029 3030 3031 3032 3033 3034 3035 3036 3037 3038 3039 3040 3041 3042 3043 3044 3045 3046 3047 3048 3049 3050 3051 3052 3053 3054 3055 3056 3057 3058 3059 3060 3061 3062 3063 3064 3065 3066 3067 3068 3069 3070 3071 3072 3073 3074 3075 3076 3077 3078 3079 3080 3081 3082 3083 3084 3085 3086 3087 3088 3089 3090 3091 3092 3093 3094 3095 3096 3097 3098 3099 3100 3101 3102 3103 3104 3105 3106 3107 3108 3109 3110 3111 3112 3113 3114 3115 3116 3117 3118 3119 3120 3121 3122 3123 3124 3125 3126 3127 3128 3129 3130 3131 3132 3133 3134 3135 3136 3137 3138 3139 3140 3141 3142 3143 3144 3145 3146 3147 3148 3149 3150 3151 3152 3153 3154 3155 3156 3157 3158 3159 3160 3161 3162 3163 3164 3165 3166 3167 3168 3169 3170 3171 3172 3173 3174 3175 3176 3177 3178 3179 3180 3181 3182 3183 3184 3185 3186 3187 3188 3189 3190 3191 3192 3193 3194 3195 3196 3197 3198 3199 3200 3201 3202 3203 3204 3205 3206 3207 3208 3209 3210 3211 3212 3213 3214 3215 3216 3217 3218 3219 3220 3221 3222 3223 3224 3225 3226 3227 3228 3229 3230 3231 3232 3233 3234 3235 3236 3237 3238 3239 3240 3241 3242 3243 3244 3245 3246 3247 3248 3249 3250 3251 3252 3253 3254 3255 3256 3257 3258 3259 3260 3261 3262 3263 3264 3265 3266 3267 3268 3269 3270 3271 3272 3273 3274 3275 3276 3277 3278 3279 3280 3281 3282 3283 3284 3285 3286 3287 3288 3289 3290 3291 3292 3293 3294 3295 3296 3297 3298 3299 3300 3301 3302 3303 3304 3305 3306 3307 3308 3309 3310 3311 3312 3313 3314 3315 3316 3317 3318 3319 3320 3321 3322 3323 3324 3325 3326 3327 3328 3329 3330 3331 3332 3333 3334 3335 3336 3337 3338 3339 3340 3341 3342 3343 3344 3345 3346 3347 3348 3349 3350 3351 3352 3353 3354 3355 3356 3357 3358 3359 3360 3361 3362 3363 3364 3365 3366 3367 3368 3369 3370 3371 3372 3373 3374 3375 3376 3377 3378 3379 3380 3381 3382 3383 3384 3385 3386 3387 3388 3389 3390 3391 3392 3393 3394 3395 3396 3397 3398 3399 3400 3401 3402 3403 3404 3405 3406 3407 3408 3409 3410 3411 3412 3413 3414 3415 3416 3417 3418 3419 3420 3421 3422 3423 3424 3425 3426 3427 3428 3429 3430 3431 3432 3433 3434 3435 3436 3437 3438 3439 3440 3441 3442 3443 3444 3445 3446 3447 3448 3449 3450 3451 3452 3453 3454 3455 3456 3457 3458 3459 3460 3461 3462 3463 3464 3465 3466 3467 3468 3469 3470 3471 3472 3473 3474 3475 3476 3477 3478 3479 3480 3481 3482 3483 3484 3485 3486 3487 3488 3489 3490 3491 3492 3493 3494 3495 3496 3497 3498 3499 3500 3501 3502 3503 3504 3505 3506 3507 3508 3509 3510 3511 3512 3513 3514 3515 3516 3517 3518 3519 3520 3521 3522 3523 3524 3525 3526 3527 3528 3529 3530 3531 3532 3533 3534 3535 3536 3537 3538 3539 3540 3541 3542 3543 3544 3545 3546 3547 3548 3549 3550 3551 3552 3553 3554 3555 3556 3557 3558 3559 3560 3561 3562 3563 3564 3565 3566 3567 3568 3569 3570 3571 3572 3573 3574 3575 3576 3577 3578 3579 3580 3581 3582 3583 3584 3585 3586 3587 3588 3589 3590 3591 3592 3593 3594 3595 3596 3597 3598 3599 3600 3601 3602 3603 3604 3605 3606 3607 3608 3609 3610 3611 3612 3613 3614 3615 3616 3617 3618 3619 3620 3621 3622 3623 3624 3625 3626 3627 3628 3629 3630 3631 3632 3633 3634 3635 3636 3637 3638 3639 3640 3641 3642 3643 3644 3645 3646 3647 3648 3649 3650 3651 3652 3653 3654 3655 3656 3657 3658 3659 3660 3661 3662 3663 3664 3665 3666 3667 3668 3669 3670 3671 3672 3673 3674 3675 3676 3677 3678 3679 3680 3681 3682 3683 3684 3685 3686 3687 3688 3689 3690 3691 3692 3693 3694 3695 3696 3697 3698 3699 3700 3701 3702 3703 3704 3705 3706 3707 3708 3709 3710 3711 3712 3713 3714 3715 3716 3717 3718 3719 3720 3721 3722 3723 3724 3725 3726 3727 3728 3729 3730 3731 3732 3733 3734 3735 3736 3737 3738 3739 3740 3741 3742 3743 3744 3745 3746 3747 3748 3749 3750 3751 3752 3753 3754 3755 3756 3757 3758 3759 3760 3761 3762 3763 3764 3765 3766 3767 3768 3769 3770 3771 3772 3773 3774 3775 3776 3777 3778 3779 3780 3781 3782 3783 3784 3785 3786 3787 3788 3789 3790 3791 3792 3793 3794 3795 3796 3797 3798 3799 3800 3801 3802 3803 3804 3805 3806 3807 3808 3809 3810 3811 3812 3813 3814 3815 3816 3817 3818 3819 3820 3821 3822 3823 3824 3825 3826 3827 3828 3829 3830 3831 3832 3833 3834 3835 3836 3837 3838 3839 3840 3841 3842 3843 3844 3845 3846 3847 3848 3849 3850 3851 3852 3853 3854 3855 3856 3857 3858 3859 3860 3861 3862 3863 3864 3865 3866 3867 3868 3869 3870 3871 3872 3873 3874 3875 3876 3877 3878 3879 3880 3881 3882 3883 3884 3885 3886 3887 3888 3889 3890 3891 3892 3893 3894 3895 3896 3897 3898 3899 3900 3901 3902 3903 3904 3905 3906 3907 3908 3909 3910 3911 3912 3913 3914 3915 3916 3917 3918 3919 3920 3921 3922 3923 3924 3925 3926 3927 3928 3929 3930 3931 3932 3933 3934 3935 3936 3937 3938 3939 3940 3941 3942 3943 3944 3945 3946 3947 3948 3949 3950 3951 3952 3953 3954 3955 3956 3957 3958 3959 3960 3961 3962 3963 3964 3965 3966 3967 3968 3969 3970 3971 3972 3973 3974 3975 3976 3977 3978 3979 3980 3981 3982 3983 3984 3985 3986 3987 3988 3989 3990 3991 3992 3993 3994 3995 3996 3997 3998 3999 4000 4001 4002 4003 4004 4005 4006 4007 4008 4009 4010 4011 4012 4013 4014 4015 4016 4017 4018 4019 4020 4021 4022 4023 4024 4025 4026 4027 4028 4029 4030 4031 4032 4033 4034 4035 4036 4037 4038 4039 4040 4041 4042 4043 4044 4045 4046 4047 4048 4049 4050 4051 4052 4053 4054 4055 4056 4057 4058 4059 4060 4061 4062 4063 4064 4065 4066 4067 4068 4069 4070 4071 4072 4073 4074 4075 4076 4077 4078 4079 4080 4081 4082 4083 4084 4085 4086 4087 4088 4089 4090 4091 4092 4093 4094 4095 4096 4097 4098 4099 4100 4101 4102 4103 4104 4105 4106 4107 4108 4109 4110 4111 4112 4113 4114 4115 4116 4117 4118 4119 4120 4121 4122 4123 4124 4125 4126 4127 4128 4129 4130 4131 4132 4133 4134 4135 4136 4137 4138 4139 4140 4141 4142 4143 4144 4145 4146 4147 4148 4149 4150 4151 4152 4153 4154 4155 4156 4157 4158 4159 4160 4161 4162 4163 4164 4165 4166 4167 4168 4169 4170 4171 4172 4173 4174 4175 4176 4177 4178 4179 4180 4181 4182 4183 4184 4185 4186 4187 4188 4189 4190 4191 4192 4193 4194 4195 4196 4197 4198 4199 4200 4201 4202 4203 4204 4205 4206 4207 4208 4209 4210 4211 4212 4213 4214 4215 4216 4217 4218 4219 4220 4221 4222 4223 4224 4225 4226 4227 4228 4229 4230 4231 4232 4233 4234 4235 4236 4237 4238 4239 4240 4241 4242 4243 4244 4245 4246 4247 4248 4249 4250 4251 4252 4253 4254 4255 4256 4257 4258 4259 4260 4261 4262 4263 4264 4265 4266 4267 4268 4269 4270 4271 4272 4273 4274 4275 4276 4277 4278 4279 4280 4281 4282 4283 4284 4285 4286 4287 4288 4289 4290 4291 4292 4293 4294 4295 4296 429

UL'YANOV, N.S.

"Extraction of Glauconite and Phosphorite from the Tailings of Concentrating plants of the Egor'yev and Upper Kama Deposits,"  
N.S. Ul'yanov, Obogashcheniye Fosforitov, Glaukonitov i Sernykh Rud, Sbornik Rabot Nauch Inst Ubobreniyam i Insektofungisidam im Ya. V. Samoylov, 1939, No 150, pp 152-66; Khim Referat Zhur 1940, No 6, pp 86 (SEE: Inst. Insect/Fungi. in Ya. V. Samoylov)

SO: U-237/49, 8 April 1949

UL'YANOV, N. S.

"Flotation of Phosphorite Ore of the Portland Horizon of the  
Egor'yev Ore Deposits," -N. S. Ul'yanov, and V. M. Ezuchevskaya,  
(Above Periodical) pp 96-103, Khim Referat Khur 1940, No 6,  
pp 84-5 (SEE: Inst. Insect/Fungi. in Ya. V. Samoylov)

SO: U-237/49, 8 April 1949

UL'YANOV, N. S.

"Flotation of the Upper Kama Phosphorite Ore Deposits," N. S.  
Ul'yanov, Above Periodical pp 103-20; Khim ~~Referat~~ Zhur, 1940, No  
6, 85 pp (SEE: Inst. Insect/Fungi. in Ya. V. Samoylov)

SO: U-237/49, 3 April 1949

UL'YANOV, N. S.

"Flotation of the Upper Kama Phosphorite Ores with a Preliminary Roasting," N. S. Ul'yanov, V. M. Vidonov, and V. G. Konstantimov, (Above Periodical) pp 120-132; Khim Referat Zhur 1940, No 6, pp 85 (SEE: Inst. Insect/Fungi. in Ya. V. Samoylov)

SO: U-237/49, 8 April 1949



UL'YANOV, N. S.

"Flotation of Phosphorite Ore from the Shchigrov Deposits,"  
N. S. Ul'yanov, and V. M. Ezuchevskaya, (Above Periodical) pp 145-51,  
Khim Referat Zhur 1940, No, 6, pp 85-6 (SEE: Inst. Insect/  
Fungi. in Ya. V. Samoylov)

SO: U-237/49, 8 April 1949

UL'YANOV, N. S.

"Flotation on a Semiproduction Scale, with Preliminary Roasting of the Ryazan-Akvilon Ore from the Egor'yev Ore Deposits," N. S. Ul'yanov, V. M. Vidonov, and N. V. Makarenko, (Above Periodical) pp 59-73, Khim Referat Zhur 1940, No 6 pp 84 (SEE: Inst. Insect/Fungi. in Ya. V. Samoylov)

SO: U-237/49, 8 April 1949

USSR/Chemistry ~~Fertilizers~~

FD-3000

Card 1/1      Pub. 50-1/17

Author      : Ul'yanov, N. S. \*

Title      : The most immediate tasks of the mined chemical raw materials industry

Periodical   : Khim. prom. No 6, 321-324, Sep 1955

Abstract    : Discusses the mining of phosphate and potassium minerals, suggesting improvements. On the basis of USA and German experience, recommends enrichment of potassium salts by flotation and expresses the opinion that the use of a hydrocyclone in combination with flotation methods is advisable. States that the gravitational method for the enrichment of Chulak-Tau and Ak-Say phosphorites is still in need of improvement, while enrichment of phosphorites by flotation has yielded good results. Says that research on the replacement of the autoclave method of melting out sulfur has lagged and should be expedited.

Institution   : Main Administration of the Mined Chemical Raw Materials Industry (\*Chief)

**"APPROVED FOR RELEASE: 03/14/2001**

**CIA-RDP86-00513R001857920015-3**

**APPROVED FOR RELEASE: 03/14/2001**

**CIA-RDP86-00513R001857920015-3"**

UL'YANOV, N.S.

Phosphate raw material and potassium fertilizers. Khim.prom.  
no.7: 430-432 O-N '57. (MIRA 10:12)  
(Phosphates) (Potassium salts)

UL'YANOV, N.S.

Conference on problems of the development of the potash industry.  
Khim. prom. no.1:54-55 Ja-F '58. (MIRA 11:3)  
(Potash industry--Congresses)



UL'YANOV, Nikolay Yegorovich; LISTOV, I.V., red.; MEL'NIKOV, V.I.,  
tekh. red.

[Outstanding people of Luzino] Znatnye liudi Luzino. Omsk,  
Omskoe knizhnoe izdatel'stvo, 1960. 70 p. (MIRA 14:12)  
(Ul'yanovskii District (Omsk Province)—Agricultural workers)



LEKAYE, V.M.; YELKIN, L.N.; UL'YANOV, N.S., kand. tekhn. nauk,  
red.

[Modern methods of sulfur recovery from sulfur ores]  
Sovremennyye sposoby polucheniia sery iz sernykh rud;  
uchebnoe posobie. Moskva, Mosk. khimiko-tekhnolog. in-t im.  
D.I. Mendeleeva, 1961. 75 p. (MIRA 16:10)  
(Sulfur)

UL'YANOV, N.S.

Problems in the development of mining, ore dressing, and chemical processing industries. Gor. zhur. no.5:3-5 ~~no~~ '63. (MIRA 16:5)

1. Gosudarstvennyy komitet po khimii pri Gosplane SSSR.  
(Apatite) (Phosphates) (Potassium) (Sulfur)

CITED SOURCE: Nauchn. tr. vuzov Dovol'naya ty'p. 1 1963, 225-233

THE PROCESS OF COMPENSATION AT THE USUALLY ENCOUNTERED 4 C. TO 10 DEGREES MUCH  
FASTER AND AT LESSON ERROR LEVELS WHEN A PHASE SENSITIVE GALVANOMETER IS EMPLOYED. 7

The comparative error introduced by the sensitive unit in such cases will be only 1.5m



UL'YANOV, O.I.

Designing a ferrodynamic galvanometer. Izv.vys.ucheb.zav.; prib.  
7 no.2:46-52 '64. (MIRA 18:4)

1. Kuybyshevskiy politekhnicheskij institut imeni Kuybysheva.  
Rekomendovana kafedroy izmeritel'ncy tekhniki.

UL'YANOV, P., polkovnik.

The eastern Pomeranian operation. Voen.snan. 29 no.9:10-11 S '53.

(MLRA 6:12)

(World War, 1939-1945--Campaigns)

UL'YANOV, P. (Astrakhan')

Cost accounting courses for radio operators. Radio no.8:40 Ag '56.  
(Astrakhan Province--Radio--Study and teaching) (MLBA 9:10)



ABRAMOV, A.A., redaktor; BOLTYANSKIY, V.G., redaktor; VASIL'YEV, A.M., redaktor; MEDVEDEV, B.V., redaktor; MYSHKIS, A.D., redaktor; NIKOL'SKIY, S.M., otvetstvennyy redaktor; POSTNIKOV, A.G., redaktor; PROKHOROV, Yu.V., redaktor; RYBNIKOV, K.A., redaktor; UL'YANOV, P.L., redaktor; USPENSKIY, V.A., redaktor; CHETAYEV, N.G., redaktor; SHILOV, G.Ye., redaktor; SHIRSHOV, A.I., redaktor; SIMKINA, Ye.N., tekhnicheskikh redaktor

[Proceedings of the third All-Union mathematical congress] Trudy tret'ego vsesoiuznogo matematicheskogo s"yezda. Moskva, Izd-vo Akademii nauk SSSR. Vol.1. [Reports of the sections] Sektsionnye doklady. 1956. 236 p. (MLRA 9:7)

1. Vsesoyuznyy matematicheskiy s"yezd. 3rd Moscow, 1956. (Mathematics)

BEREZOVIKO, P.; KOZHEVNIKOV, N., inzh.-tekhnolog; M.L'NIKOV, A.;  
UL'YANOV, P., konditer

Advice to the cook. Obshchestv.pit. no.11:16-17 N '59.  
(MIRA 13:3)

1. Upravleniye rabocheho snabzheniya Sverdlovskogo sovnarkhoza  
(for Kozhevnikov).

(Cookery)

UL'YANOV, P.

Party organization of the interfarm building organizations.  
Sel'.stroï. 15 no.8:12-14 Ag '60. (MIRA 13:8)

1. Sekretar' partorganizatsii Gul'kevichskogo meshkolkhozstroya  
Krasnodarskogo kraya.  
(Krasnodar Territory--Building)  
(Collective farms--Interfarm cooperation)

UL'YANOV, P., kand.economicheskikh nauk

Socialist economy is the indestructible basis for our country's defenses.  
Ty1 1 snab.Sov.Voor.311 21 no.2:10-15 F '61. (MIRA 14:6)  
(Russia—Economic conditions)

UL'YANOV, P., kand.ekonomicheskikh nauk

Communism is an abundance. Komm.Vooruzh.Sil 2 no.3:39-47 F '62.  
(MIRA 15:1)

(Cost and standard of living)

UKYANOV, P. A.

AID Nr 972-17 20 May

VACUUM CLADDING OF REFRACTORY METALS (USSR)

Ul'yanov, P. A., N. D. Tarasov, and S. F. Koftun. Tsvetnyye metally, no. 3, Mar 1963, 74-76. S/136/63/000/003/003/004

The cladding of Nb, Mo, and Ta with 1X18H9T [AISI-321] stainless steel, Ni-chrome, Zh-602 alloy [3% Fe, 0.35-0.75% Al and Ti, 0.4% Mn, 19-22% Cr, 1.8-2.3% Mo, 0.8% Si, 0.08% C, 1.3-1.8% Nb], and zirconium has been investigated experimentally. Cladding was performed in a vacuum rolling mill designed by the Physicotechnical Institute of the Ukrainian Academy of Sciences. Refractory billets were mechanically cleaned or pickled, spot welded or riveted to the cladding material, heated in vacuum to the rolling temperature, and then rolled to the required thickness. Pressure in the vacuum system during heating and rolling was maintained at  $4 \cdot 10^{-5}$  mm Hg or lower. In order to prevent work hardening, the rolling temperature was maintained above that of the recrystallization of the rolled metal. The strength of the

Card 1/2

AID Nr. 971-17 20 May

VACUUM CLADDING [Cont'd]

S/136/63/000/003/004

bond between the cladding and the base metal was found to increase with increasing reduction and with higher rolling temperatures. Microhardness tests showed that Mo and Cr-Ni alloy claddings do not form chemical compounds in the interface zone; A sharp increase of interface microhardness from ~ 230 to 740 kg/mm<sup>2</sup> was observed in Nb clad with 3M-602 alloy. Some hardness increase was observed in Nb clad with Zr or Ti. Aging at 1200°C for 2 hrs had little or no effect on the structure or strength of the bond between Mo or Nb and Cr-Ni alloy cladding; aging at 1200°C for 10 hrs increased bond strength by 15-20%. Shear strength of the bond between niobium and zirconium cladding rolled at 1100°C with reductions of 20 or 40% was ~ 30 or 64 kg/mm<sup>2</sup>, respectively, and that between molybdenum and 3M-602 cladding rolled at 1190°C with reductions of 20 or 45% was ~ 28 or 43 kg/mm<sup>2</sup>, respectively.

[AZ]

Card 2/2

UL'YANOV, P.L.

Series in Haar's system. Vest. Mosk. un. Ser. 1: Mat., mekh.  
20 no.4:35-43. J1-Ag '65. (MIRA 18:9)

1. Kafedra teorii funktsii i funktsional'nogo analiza Moskovskogo  
gosudarstvennogo universiteta imeni M.V. Lomonosova.



ULYANOV, T. L.

## USSR.

Ul'yanov, P. L. On some equivalent conditions of convergence of series and integrals. Uspehi Matem. Nauk (N.S.) 3 no. 6:58, 133-141 (1953). (Russian)

1 - F/W

(N. S. 13 no. 6:136), 133-141 (1953). (Russian.)

Suppose  $f \in L_{p, \omega}(T)$  and has the Fourier coefficients  $f_k$ . Let  $\omega(t)$  be a function in the class of the integral  $\int_0^t f(x+t) f(x-t) dx$  and the series  $\sum_{k=0}^{\infty} \omega(k) a_k^2$ , where the function  $a(t)$  and the sequence  $\omega(k)$  depend on each other (given  $t$  or  $\omega(k)$ ), satisfying the conditions:

1.  $\omega(k) \geq 0$  and  $\omega(k)$  decreases slowly to 0 as  $k \rightarrow \infty$ .
2.  $\omega(k) \rightarrow 0$  as  $k \rightarrow \infty$  and  $\omega(k) \rightarrow 0$  as  $k \rightarrow \infty$  for which  $\delta(f, \omega) = 0$  or  $\delta(f, \omega) = \infty$  versus  $\omega(k)$  is chosen as  $f_k^2 \omega(k)$  or  $1/k^2 \omega(k)$ , respectively. In the case  $\omega(t) = 1, \omega(k) = 1$  and the equivalence reduces to the well known theorem of Plessner. In some cases when the sequence  $\omega(k)$  increases too rapidly the finiteness of the series is equivalent to that of the integral obtained by replacing  $|f(x+t) - f(x-t)|$  by a higher difference.

G. Klein (South Hadley, Mass.).

G. Klein (South Hadley, Mass.).

following Plossner, the so obtained criterion is shown equivalent to the condition  $\sum_{k=1}^{\infty} (a_k^2 + b_k^2) \log k < +\infty$  of Kolmogorov-Schilverstov as applied to the product of  $f$  and the

Mathematical Reviews  
Vol. 15 No.1  
Jan. 1954  
Analysis

7-13-54  
LL

Ul'yanov, P. L. On trigonometric series with monotonically decreasing coefficients. Doklady Akad. Nauk SSSR (N.S.) 90, 33-36 (1953). (Russian)

The author considers the functions  $f(x) = \sum_{k=1}^{\infty} a_k \cos kx$ ,  $f(x) = \sum_{k=1}^{\infty} a_k \sin kx$ , under the condition that  $a_k \rightarrow 0$  and  $\sum |\Delta a_n| < \infty$ . It is well known that neither series need be a Fourier series under this condition, if integration is Lebesgue integration. The author says that a measurable function  $\phi(x)$  is A-integrable on  $(a, b)$  if the measure of the set where  $|\phi(x)| \geq n$  is  $o(1/n)$  and the Lebesgue integral of the function obtained by truncating  $\phi(x)$  at  $\pm n$  approaches a limit. [See, e.g., Očn. Mat. Sbornik, N.S. 28(70), 293-336 (1951); these Rev. 13, 20.] The following theorems are given. (1) The series for  $f(x)$  and  $f(x)$  are the A-Fourier series of their sums. (2) If  $\phi(x)$  is a function of bounded variation whose conjugate  $\bar{\phi}(x)$  is also of bounded variation,  $(A)f_{\sigma}^{\tau} f \bar{\phi} = -(A)f_{\sigma}^{\tau} f \phi$ . (3) If  $\phi(x)$  is a bounded measurable function, the definition of  $\bar{\phi}(x)$  as a Cauchy integral agrees almost everywhere with the definition of  $\bar{\phi}(x)$  as an A-integral. (4) For all  $x$ ,  $f(x)$  is the negative of the A-conjugate of  $f(x)$ ; except perhaps at  $0, \pm\pi$ ,  $f(x)$  is the A-conjugate of  $f(x)$ .

R. P. Boas, Jr. (Ulan-Ude, U.S.S.R.).

UL'YANOV, P. L.  
USSR/Mathematics - Fourier series

FD-1427

Card 1/1 : Pub. 64 - 5/9

Author : Ul'yanov, P. L. (Moscow)

Title : Application of A-integration to a class of trigonometric series

Periodical : Mat. sbor., 35 (77), pp 469-490, Nov-Dec 1954

Abstract : The main results of this work were formulated without proof in the author's article "Trigonometric series with monotonically decreasing coefficients. "DAN SSSR, 90, No 1, 33-36, 1953. In the present work the author gives the principal definitions and cites certain works devoted to the same problem. He proves that  $f(x) = a_0 + \sum a_k \cos kx$  is a Fourier (A) series of  $f(x)$  and its sine-conjugate  $f(x)$ . Thirteen references, 2 USSR.

Institution : .

Submitted : October 28, 1953

*Ulyanov P.L.*

*MS* ✓ Ulyanov, P. L. Some questions of  $A$ -integration. Dokl. Akad. Nauk SSSR (N.S.) 102 (1955), 1977-1980. 1- F/W

(Russian)  
A measurable real-valued function  $f$  on  $[a, b]$  is said to be  $A$ -integrable if

$$(1) \quad m\{x: x \in [a, b], |f(x)| < n\} = o(n^{-1})$$

and

$$(2) \quad \lim_{n \rightarrow \infty} \int_a^b \min[\max(f(x), -n), n] dx = (A) \int_a^b f(x) dx$$

exists and is finite. This notion is attributed to Kolmogorov, and differs hardly at all from the  $Q$ -integral of Fitchmarsh [Proc. London Math. Soc. (2) 29 (1928), 49-80]. [For other applications of this notion, see Ulyanov same Dokl. (N.S.) 90 (1953), 33-36; Mat. Sb. N.S. 35(77) (1954), 469-490; MR 15, 27; 16, 467.] The author states that Kolmogorov proved property (1) for all functions on  $[0, 2\pi]$  conjugate to functions in  $L_1(0, 2\pi)$  [Fund. Math. 7 (1925), 24-29], but Kolmogorov seems only to have proved that the left side is  $o(n^{-1})$ . For  $f \in L_1(0, 2\pi)$ , let  $\bar{f}$  denote the conjugate function of  $f$ . Theorem: If  $f \in L_1(0, 2\pi)$  and if  $g$  and  $\bar{g}$  are essentially bounded, then

$1/2$  ✓

$\frac{1}{\text{(over)}}$

Uhlenbrock, P.L.

$$(A) \int_0^{2\pi} f(x)g(x)dx = - \int_0^{2\pi} f(x)g'(x)dx,$$

Theorem: If  $f \in L_p(0, 2\pi)$  ( $p > 1$ ) and  $f$  has period  $2\pi$ , then

$$(A) \int_0^{2\pi} [f(x+t) - f(x-t)] \frac{1}{2} \cot \frac{1}{2}t dt = -\pi f(x)$$

for almost all  $x \in [0, 2\pi]$ . A formula is also given for inverting the transform  $f \rightarrow F$  ( $f \in L_1(-\pi, \pi)$ ). Proofs are sketched.

E. Hewitt (Princeton, N.J.).

2/2

Small

ULJANOV, P.L.  
 SUBJECT USSR/MATHEMATICS/Theory of functions  
 AUTHOR ULJANOV P.L.  
 TITLE On the continuation of functions.  
 PERIODICAL Doklady Akad. Nauk 105, 913-915 (1955)  
 reviewed 7/1956

CARD 1/2

PG -- 182

The author considers a function  $f(x)$  which is defined on  $[\alpha, \beta]$  and on  $[a, b] \subseteq [\alpha, \beta]$  has the property A. He seeks a function  $f_1(x)$  which is defined on  $[c, d]$  (where  $(c, d) \supset [a, b]$ ), on  $[a, b]$  identical with  $f(x)$  and on  $[c, d]$  possesses the property A. Beside of  $f(x)$  its conjugate function

$$\bar{f}(x) = - \lim_{\varepsilon \rightarrow 0} \frac{1}{\pi} \int_{\varepsilon}^{\pi} \frac{f(x+t) - f(x-t)}{2 \operatorname{tg} \frac{1}{2} t} dt$$

is considered.

For integrable and continuous functions the following theorems are formulated and a sketchy proof is given: 1. Let the periodic function  $f(x) \in L(0, 2\pi)$  have the property that  $f(x)$  and  $\bar{f}(x)$  are integrable on  $[a, b] \subset [0, 2\pi]$  and for a  $\varepsilon > 0$  holds:

$$\int_0^n f(b+t) dt = O \left\{ \left( \ln \frac{1}{|n|} \right)^{-1-\varepsilon} \right\}, \quad \int_0^n f(a+t) dt = O \left\{ \left( \ln \frac{1}{|n|} \right)^{-1-\varepsilon} \right\}.$$

Then there exists a function  $\varphi(x)$  such that  $\varphi(x) = f(x)$  on  $[a, b]$  and  $\varphi(x) \in L(0, 2\pi)$ ,  $\bar{\varphi}(x) \in L(0, 2\pi)$ . 2. Let  $f(x) \in L(0, 2\pi)$  be periodic,  $f(x)$  and

Doklady Akad.Nauk 105, 913-915 (1955)

OARD 2/2

PQ - 182

$\bar{f}(x)$  continuous on  $[a, b] \subset [0, 2\pi]$ . Then  $f(x)$  can be continued from  $[a, b]$  to  $[0, 2\pi]$  such that it and its conjugate function are continuous on the whole interval  $[0, 2\pi]$ . 3. Let  $f(x) \in L(0, 2\pi)$  be periodic,  $f(x)$  and  $\bar{f}(x)$  essentially bounded on  $[a, b] \subset [0, 2\pi]$  and

$$\int_0^t f(a+n)dn = O(|t|), \quad \int_0^t f(b+n)dn = O(|t|)$$

$$\lim_{h \rightarrow 0} \left| \int_h^\pi \frac{f(a+n)-f(a-n)}{n} dn \right| < \infty, \quad \lim_{h \rightarrow 0} \left| \int_h^\pi \frac{f(b+n)-f(b-n)}{n} dn \right| < \infty,$$

then  $f(x)$  can be continued from  $[a, b]$  to  $[0, 2\pi]$  such that the property of the essential boundedness for  $f$  and  $\bar{f}$  remains true.

INSTITUTION: Lomonossov University, Moscow



ABRAMOV, A.A., redaktor; BOITYANSKIY, V.G., redaktor; VASIL'YEV, A.M., redaktor; MEDVEDEV, B.V., redaktor; MYSHKIS, A.D., redaktor; NIKOL'SKIY, S.M., otvetstvennyy redaktor; POSTNIKOV, A.G., redaktor; PROKHOROV, Yu.V., redaktor; RYBNIKOV, K.A., redaktor; UL'YANOV, P.L., redaktor; USPENSKIY, V.A., redaktor; GHETAYEV, N.G., redaktor; SHILOV, G.Ye., redaktor; SHIRSHOV, A.I., redaktor; SIMKINA, Ye.H., tekhnicheskiy redaktor

[Proceedings of the all-Union Mathematical Congress] Trudy tret'ego vsesoiuznogo Matematicheskogo s'ezda; Moskva i iun'-iul' 1956. Moskva, Izd-vo Akademii nauk SSSR. Vol.2. [Brief summaries of reports] Kratkoe sodержanie obzornykh i sektiionnykh dokladov. 1956. 166 p. (MLRA 9:9)

1. Vsesoyuznyy matematicheskiy s'yezd. 3, Moscow, 1956. (Mathematics)

*U.L. YAGLOV, P.L.*

1-FW

*Ulyanov, P. L. On the A-Cauchy Integral. I. Uspehi  
Mat. Nauk (N.S.) 11 (1956), no. 5(71), 223-229. (Rus-  
sian)*

The A-integral of  $\phi(x)$  is the limit of the L-integral of  
the function obtained by truncating  $\phi(x)$  at  $\pm n$  where  
the measure of the set where  $|\phi(x)| \geq n$  is  $O(1/n)$ . The author  
shows that if a real  $f$  belongs to  $L$  on the interval

where the Cauchy integral, it is represented by the A-  
Cauchy integral of its boundary values. Various corollaries  
are obtained. R. P. Boas, Jr. (Evanston, Ill.)

SUBJECT USSR/MATHEMATICS/Fourier series CARD 1/2 PG - 742  
 AUTHOR ULJANOV P.L.  
 TITLE On almost everywhere permanently convergent series.  
 PERIODICAL Mat.Sbornik,n.Ser. 40, 1, 95-100 (1956)  
 reviewed 5/1957

An almost everywhere permanently convergent function series is a series which converges almost everywhere for an arbitrary transposition of the terms.

Let  $\{P_n(x)\}$  ( $n=0,1,2,\dots$ ) be a system of polynomials, being defined on  $[a,b]$ , being complete with respect to  $L$  and closed with respect to  $L^2$ , which is orthonormalized with the weight  $\tau(x)$  ( $\tau(x)$  is defined on  $[a,b]$ , positive and integrable). The series

$$(1) \quad \sum_{k=0}^{\infty} c_k P_k(x)$$

is called the Fourier series of the integrable function  $f(x)$  if

$$c_k = \int_a^b f(x) \tau(x) P_k(x) dx \quad (k=0,1,2,\dots).$$

Let  $\omega(\delta, f)$  be the modulus of continuity of  $f$  on  $[a,b]$  with the length of

Mat.Sbornik, n.Ser. 40, 95-100 (1956)

CARD 2/2

PG - 742

steps  $\delta$ . Joining the results of Kolmogorov (Doklady Akad.Nauk 1, 291-294 (1934)) and Natanson (Doklady Akad.Nauk 2, 209-211 (1934)) the author proves the theorems:

1. If  $f(x) \in L(a,b)$  and

$$\omega(\delta, f) = o \left\{ \frac{1}{\ln \frac{1}{\delta} (\ln \ln \frac{1}{\delta})^{1+\varepsilon}} \right\} \text{ for } \delta \rightarrow +0,$$

then the Fourier series (1) of the function  $f(x)$  on  $[a,b]$  converges almost everywhere for an arbitrary arrangement of the terms.

2. If  $f(x)$  is of bounded variation on  $a,b$  and if

$$0 < \tau(x) \leq \frac{\Delta}{\sqrt{(b-x)(x-a)}} \text{ for } x \in [a,b],$$

then for every  $\varepsilon > 0$  there holds

$$\sum_{k=0}^{\infty} |c_k|^{1+\varepsilon} < +\infty \quad \sum_{k=0}^{\infty} c_k^2 k^{1-\varepsilon} < +\infty.$$

INSTITUTION: Moscow.

UL'YANOV, P.L.

A-integral and conjugate functions. Uch. zap. Mosk. un. no.181:  
139-157 '56. (MLRA 10:4)  
(Fourier's series) (Integrals)

The author proves the theorems announced in Uspehi  
Mat. Nauk (N.S.) 10 (1955), no. 1(63), 189-191.  
*R. P. Boas, Jr. (Evanston, Ill.).*

UL'YANOV, P. L.

Call Nr: AF 1108825

r Transactions of the Third All-union Mathematical Congress (Cont.) Moscow, Jun-Jul '56, Trudy '56, V. 1, Sect. Rpts., Izdatel'stvo AN SSSR, Moscow, 1956, 237 pp. There are 6 references, all of them USSR.

UL'yanov, P. L. (Moscow). About A-integrals of Cauchy. 107-108

Fedorov, V. S. (Ivanovo). On Monogenic Functions. 108-109

Fishman, K. M. (Chernovitsy). On a Class of Hilbert Spaces of Analytic Function. 109

Fuksman, N. A. (Tashkent). About Analytic Functions of Integral Complex Argument. 109-110

Mention is made of Romanov, N. P.

Khavinson, S. Ya. (Moscow). P. L. Chebyshev's Systems and the Uniqueness of the Best Polynomial Approximation in the Metrics of  $L_1$  Space. 110

Card 34/80

ULYANOV, P.L.

SUBJECT  
AUTHOR  
TITLE  
PERIODICAL

USSR/MATHEMATICS/Theory of functions  
ULJANOV P.L.  
On Cauchy  $\Lambda$ -integrals on curves.  
Doklady Akad. Nauk 112, 383-385 (1957)  
reviewed 4/1957

CARD 1/3

PG - 724

In the complex  $\zeta$ -plane let be given a smooth curve  $l$  of the length  $l_0$  beginning in  $\zeta_0$  and ending in  $\zeta'_0$ . Its equation be  $\zeta = \tau(s) = \tau_1(s) + i\tau_2(s)$ , where  $s$  is the arc length of  $\zeta_0$  to  $\zeta$  ( $\zeta_0 = \tau(0)$ ,  $\zeta'_0 = \tau(l_0)$ ). Then the function  $f(\zeta) = f_1(s) + if_2(s)$  being defined on  $l$  is called  $\Lambda$ -integrable on  $l$  if the functions

$$\varphi_1(s) = [f_1(s)\tau'_1(s) - f_2(s)\tau'_2(s)]$$

$$\varphi_2(s) = [f_2(s)\tau'_1(s) + f_1(s)\tau'_2(s)]$$

are  $\Lambda$ -integrable on the line  $0 \leq s \leq l_0$  (as to the  $\Lambda$ -integrability on lines compare Titchmarsh, Proc.London Math.Soc. 29, 49 (1929)). The complex number



Doklady Akad.Nauk 112, 383-385 (1957)

CARD 2/3

PG - 724

$$I = (\Delta) \int_0^1 \varphi_1(s) ds + i(\Delta) \int_0^1 \varphi_2(s) ds$$

is called the  $\Delta$ -integral of the function  $f(\zeta)$  on the curve  $l$

$$(\Delta) \int_l f(\zeta) d\zeta = I.$$

With the aid of this definition the following principal result can be formulated: Let  $l$  be a closed curve which limits the domain  $G$ . Its equation be  $\zeta = \zeta(s) = x(s) + iy(s)$ , where

$$|x'(s_2) - x'(s_1)| \leq k |s_2 - s_1|^\alpha, \quad |y'(s_2) - y'(s_1)| \leq k |s_2 - s_1|^\alpha$$

for all  $s_1, s_2$  and certain constant  $k > 0, \alpha > 0$ . If the analytic function  $F(z)$  is representable in  $G$  by an  $L$ -integral of the Cauchy type, i.e. if

$$F(z) = \frac{1}{2\pi i} (L) \int_l \frac{f(\zeta)}{\zeta - z} d\zeta \quad (z \in G, f(\zeta) \in L(l)),$$

Doklady Akad.Nauk 112, 383-385 (1957)

CARD 3/3

PG - 724

then

$$F(z) = \frac{1}{2\pi i} (\Delta) \int_1 \frac{F_1(\zeta)}{\zeta - z} d\zeta,$$

where  $F_1(\zeta)$  are the limit values of the function  $F(z)$  if  $z$  coming from the interior of  $G$  reaches 1. Some conclusions are given.

20-4-12/51

UL'YANOV, P.L.

AUTHOR:

UL'YANOV, P.L.

TITLE:

On Permutations of a Trigonometric System (O perestankakh trigonometricheskoy sistemy)

PERIODICAL:

Doklady Akademii Nauk SSSR, 1957, Vol. 116, Nr. 4, pp. 568-571 (USSR)

ABSTRACT:

Let

$$(1) \quad \frac{a_0}{2} + \sum_{k=1}^{\infty} (a_k \cos kx + b_k \sin kx)$$

be the Fourier series of  $f(x) \in L(0, 2\pi)$ ,  $f(x+2\pi) = f(x)$ . (1) is called unconditionally convergent almost everywhere if it converges almost everywhere after an arbitrary permutation of the terms. Let  $E_n^{(2)}(f)$  be the best approximation of  $f(x)$  in the metric of the  $L^2$  by trigonometric polynomials of the order  $(n-1)$ .

Theorem: If  $\sum_{n=10}^{\infty} \frac{(\ln \ln n)^{1+\varepsilon} \ln n}{n} \{E_n^{(2)}(f)\}^2 < \infty$ ,  $\varepsilon > 0$ , then (1) converges unconditionally almost everywhere on  $[0, 2\pi]$ .

Theorem: If for  $\varepsilon > 0$  there holds:

$$\int_0^{2\pi} \int_0^{2\pi} \frac{|\ln t| |\ln |\ln t||^{1+\varepsilon}}{t} [f(x+t) - f(x-t)]^2 dx dt < \infty,$$

Card 1/2

On Permutations of a Trigonometric System

20-4-12/51

then (1) is unconditionally convergent almost everywhere on  $[0, 2\pi]$ .

Theorem: There exists a continuous  $2\pi$ -periodic function  $f(x)$  the Fourier series of which after a certain permutation of the terms does not converge on  $[0, 2\pi]$  for every  $q > 2$  in the metric of the  $L^q$ .

Several further similar results are given which e.g. generalize well known results due to Marcinkiewicz [Ref 3] and Orlicz [Ref.8].

ASSOCIATION: Moscow State University im. M.V. Lomonosov (Moskovskiy gosudarstvennyy universitet im. M.V. Lomonosova)

PRESENTED BY: A. N. Kolmogorov, Academician, April 10, 1957

SUBMITTED: February 28, 1957

AVAILABLE: Library of Congress

Card 2/2

KACHMAZH, S. [Kaczmarz, Stefan]; SHTINGAUZ, G.; GUTER, R.S. [translator];  
UL'YANOV, P.L. [translator]; VILENKIN, N.Ya., red.

[Theory of orthogonal series] Teoriia ortogonal'nykh riadov.  
Pod red. i s dop. N.IA.Vilenkina. Moskva, Gos.izd-vo fiziko-  
matem.lit-ry, 1958. 507 p. (MIRA 12:11)  
(Series, Orthogonal)

NIKOL'SKIY, S.M., otv.red.; ABRAMOV, A.A., red.; BOLTYANSKIY, V.G., red.;  
VASIL'YEV, A.M., red.; MEDVEDEV, B.V., red.; MYSHKIS, A.D., red.;  
POSTNIKOV, A.G., red.; PROKHOROV, Yu.V., red.; RYBNIKOV, K.A.,  
red.; UL'YANOV, P.L., red.; USPENSKIY, V.A., red.; CHETAYEV, N.G.,  
red.; SHILOV, G.Ye., red.; SHIRSHOV, A.I., red.; GUSEVA, I.N.,  
tekhn.red.

[Proceedings of the Third All-Union Mathematical Congress] Trudy  
tret'ego Vsesoiuznogo matematicheskogo s"ezda. Vol.3 [Synoptic  
papers] Obzornye doklady. Moskva, Izd-vo Akad.nauk SSSR. 1958. 596 p.  
(MIRA 12:2)

1. Vsesoyuznyy matematicheskiy s"ezd. 3d, Moscow, 1956.  
(Mathematics--Congresses)

P.L. 01/10/1957

16(1)

**AUTHORS:**

Shorin, I.A., University Lecturer, and  
Kopylov, V.D., Scientific Assistant  
Lomonosov - lectures 1957 at the Mechanical-Mathematical  
Faculty of Moscow State University (Lomonosovskie  
obshchiye 1957 goda za matematiko-matematicheskoy fakultete  
MSU)

**TITLE:**

**PERIODICAL:**

Vestnik Moskovskogo Universiteta. Seriya Matematiki, Mekhanika,  
astronomiya, fizika, khimiya, 1958, str. 241-246 (USSR)

**ABSTRACT:**

The Lomonosov lectures 1957 took place from October 17 -  
October 31, 1957 and were dedicated to the 40-th anniversary  
of the October revolution.

16. A.D. Gorbunov, Lecturer and B.M. Budak, Lecturer -  
Difference Methods for the Solution of Hyperbolic  
Equations.

17. M.S. Mikheylov - Number of Calculation Operations for  
the Solution of Elliptic Equations.

18. V.I. Lebedev, Assistant - Difference Method for the  
Solution of the Sobolev-Selivskiy System.

19. Professor Ye.B. Dyukin - Markov Processes and Semigroups.

20. A.G. Kostyuchenko, Candidate of Physical-Mathematical  
Sciences - Decomposition of Differential Operators With  
Respect to Generalized Eigenfunctions.

21. F.A. Berezin, Candidate of Physical-Mathematical Sciences,  
Foundations of the Theory of Spherical Harmonics on Mani-  
folds.

22. V.K. Borok, Aspirant - General Properties of Partial  
Evolution Systems.

23. S.A. Kuznetsov, Candidate of Physical-Mathematical  
Sciences - On Constructive Mathematical Analysis.

24. P.M. Gulyaev, Lecturer - Reversal of Terms in Trigonometric Series.

25. I.G. Petrovskiy, Academician and Ye.K. Landis, Senior  
Scientific Assistant - On the Number of Boundary Cycles  
of a Differential Equation of First Order With a Rational  
Right Side.

The contents of all the lectures have already been published.

Cart 5/5

12

16(+) 16,4100

AUTHOR: Ul'yanov, P.L.

SOV/155-58-4-11/34

TITLE: On the Divergence of Orthogonal Series to  $+\infty$  (0 raskhodimosti ortogonal'nykh ryadov k  $+\infty$ )

PERIODICAL: Nauchnyye doklady vysshey shkoly. Fiziko-matematicheskiye nauki, 1958, Nr 4, pp 63 - 68 (USSR)

ABSTRACT: Let

$a_n > 0$  and  $\sum_{n=1}^{\infty} a_n^2 = \infty$ . Then there exists a system

$\{\varphi_n(x)\}$  of bounded functions orthogonally normed on  $[0,1]$

so that the orthogonal series  $\sum_{k=1}^{\infty} b_k \varphi_k(x)$  for every order of the terms diverges everywhere on  $[0,1]$  to  $+\infty$ , if  $b_k \geq a_k$ .

Theorem :

It exists an orthogonal series  $\sum_{n=1}^{\infty} c_n \varphi_n(x)$ , which for an arbitrary sequence of the terms diverges everywhere on  $[0,1]$

Card 1/2



On the Divergence of Orthogonal Series to  $+\infty$

SOV/155-58-4-11/34

to  $+\infty$ , while  $\sum_{n=1}^{\infty} |c_n|^{2+\varepsilon} < \infty$  is for every  $\varepsilon > 0$ .

Theorem : On  $[0,1]$  there exists an orthogonal series  $\sum_{n=1}^{\infty} c_n \varphi_n(x)$

with the properties: 1.) it diverges to  $+\infty$  everywhere on  $[a,b] \subset (0,1)$  for arbitrary reversal of the terms 2.) the orthogonally normed system  $\{\varphi_n(x)\}$  is bounded on  $[0,1]$ .

The author mentions D.Ye. Men'shov.

There are 3 references, 2 of which are Soviet, and 1 French.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet imeni M.V.Lomonosova  
(Moscow State University imeni M.V. Lomonosov)

SUBMITTED: June 4, 1958

Card 2/2

SOV/38-22-4-4/6

AUTHOR: Ul'yanov, P.L.

TITLE: On the Series With Respect to a Transposed Trigonometric System (O ryadakh po perestavlennoy trigonometricheskoy sisteme)

PERIODICAL: Izvestiya Akademii nauk SSSR, Seriya matematicheskaya, 1958, Vol 22, Nr 4, pp 515-542 (USSR)

ABSTRACT: § 1. Theorem: Let  $f(x) \in L^2(0, 2\pi)$  and for an  $\varepsilon > 0$  let be

$$\sum_{n=10}^{\infty} \frac{(\ln \ln n)^{1+\varepsilon}}{n} \left\{ E_n^{(2)}(f) \right\}^2 < \infty, \text{ where } E_n^{(2)}(f) \text{ is the best}$$

approximation of  $f(x)$  in the metric  $L_2$  by trigonometric polynomials of order  $\leq n - 1$ . Then the Fourier series of  $f(x)$  converges absolutely almost everywhere on  $[0, 2\pi]$  (i.e. under arbitrary transposition of the terms). Theorem: If  $f(x) \in L^2(0, 2\pi)$  and if for an  $\varepsilon > 0$  it holds :

Card 1/ 4

On the Series With Respect to a Transposed Trigonometric System SOV/38-22-4-4/6

$$\int_0^{2\pi} \int_0^{2\pi} \frac{|\ln t| |\ln |\ln t||^{1+\varepsilon}}{t} [f(x+t) - f(x-t)]^2 dt dx < \infty, \text{ then the}$$

Fourier series of  $f(x)$  is absolutely convergent almost everywhere on  $[0, 2\pi]$ . § 2 deals with the summability of the series

$$\frac{a_0}{2} + \sum_{\nu=1}^{\infty} (a_{\nu} \cos k_{\nu} x + b_{\nu} \sin k_{\nu} x), \text{ where all } k_{\nu} \text{ are integer}$$

and different. It is shown, that even the Fourier series with respect to a transposed system also with relatively strong Töplitz methods need no longer be summable.

§ 3 Theorem: There exists a fixed transposed trigonometric system  $\{\cos m_{\nu} x, \sin m_{\nu} x\}$  with the properties 1.) For all  $1 \leq p < 2$  there exists an  $f(x) \in L^p(0, 2\pi)$  with derivatives of arbitrary order continuous on  $(0, 2\pi)$  and with  $f(x) = 0$  for  $x \in [1, 2\pi - 1]$ ; the Fourier series

$$f(x) \sim \frac{a_0}{2} + \sum_{\nu=1}^{\infty} a_{m_{\nu}} \cos m_{\nu} x + b_{m_{\nu}} \sin m_{\nu} x$$

Card 2/ 4

On the Series With Respect to a Transposed Trigonometric System SOV/38-22-4-4/6

of which diverges almost everywhere on  $[0, 2\pi]$  and does not converge in the metric  $L$ ; also the Fourier series for the conjugate function

$$\bar{f}(x) = -\frac{1}{\pi} \lim_{\epsilon \rightarrow 0} \int_{\epsilon}^{\pi} \frac{f(x+t) - f(x-t)}{2 \operatorname{tg} \frac{1}{2} t} dt \text{ diverges}$$

indefinitely on  $[0, 2\pi]$  and does not converge in the metric  $L$ . 2.) There exists a continuous function  $\varphi(x)$ , the Fourier series of which with respect to the system  $\{\cos m_p x, \sin m_p x\}$  does not converge on  $[0, 2\pi]$  in the metric  $L^p$  for any  $p > 2$ . Constructive proof. § 4 brings several conclusions; e.g. it is proved that the transposed system forms in general for  $p \in [1, 2) + (2, \infty)$  no base in  $L^p(0, 2\pi)$ . Also the Riemannian localization principle does not hold in general for the transposed system. Similar statements are given in the complex domain. Altogether there are given 27 definitions, theorems, conclusions and remarks.

Card 3/4

On the Series With Respect to a Transposed Trigonometric System SOV/38-22-4-4/6

There are 12 references, 6 of which are Soviet, and 6 Polish.

PRESENTED: by Aleksandrov, P.S., Academician

SUBMITTED: October 11, 1957

1. Mathematics 2. Trigonometry 3. Fourier's series

Card 4/4

16(1)

AUTHOR:

Ul'yanov, P.L.

SOV/38-22-6-4/6

TITLE:

On Unconditional Convergence and Summability (O bezuslovnoy skhodimosti i summiruyemosti)

PERIODICAL:

Izvestiya Akademii nauk SSSR, Seriya matematicheskaya, 1958, Vol 22, Nr 6, pp 811 - 840 (USSR)

ABSTRACT:

The author investigates the connection between the unconditional convergence and summability for trigonometric and orthogonal series. § 1 contains several auxiliary theorems, § 2 considers trigonometric series. Among others it is shown that "unconditional summability" is equivalent to an "unconditional convergence almost everywhere". Furthermore it is shown that the transposed Fourier series of the functions  $f(x) \in L^p(0, 2\pi)$  for  $p > 2$  are in general almost everywhere summable with no Toeplitz method. In § 3 it is investigated under which conditions the results of § 2 can be transferred to orthogonal series. Moreover it is tried to explain why in certain cases the results for orthogonal series deviate from those trigonometric series. 11 theorems and more than 20 lemmata, consequences, etc are brought.

Card 1/2

On Unconditional Convergence and Summability

SOV/38-22-6-4/6

There are 11 references, 5 of which are Soviet, 5 Polish, and 1 German.

PRESENTED: by S.L. Sobolev, Academician

SUBMITTED: September 29, 1957

Card 2/2

UL'YANOV, P. L., Doc Phys-Math Sci (diss) -- "A Cauchy-type integral. Convergence and summability". Moscow, 1959. 8 pp (Moscow Order of Lenin and Order of Labor Red Banner State U im M. V. Lomonosov), 150 copies (KL, No 9, 1960, 121)



67508

SOV/155-59-1-11/30

16(1) 16,4000

AUTHOR:

Ul'yanov, P.L.

TITLE:

Unconditional Convergence With Respect to  $+\infty$   
Nauchnyye doklady vysshey shkoly. Fiziko-matematicheskiye nauki,  
1959, Nr 1, pp 71 - 80 (USSR)

PERIODICAL:

ABSTRACT:

The series  $\sum_{n=1}^{\infty} f_n(x)$

$(x \in E)$

is said to be unconditionally

convergent with respect to  $+\infty$  on the set  $E$  if for an arbitrary arrangement of the terms on  $E$  it converges to  $+\infty$ .  
Basing on his earlier results the author proves six theorems on series unconditionally convergent with respect to  $+\infty$  and other connected questions.

Theorem 1 : To every sequence  $\{a_n\}$  with  $\sum_{n=1}^{\infty} a_n^2 = \infty$  there exists an orthogonal series  $\sum_{n=1}^{\infty} a_n \varphi_n(x)$  which on  $[0,1]$  is

14

67508

SOV/155-59-1-11/30

Unconditional Convergence With Respect to  $+\infty$

unconditionally convergent with respect to  $+\infty$ .

Theorem 2 : To every sequence  $\{a_n\}$  with

$$(2) \quad \sum_{n=1}^{\infty} a_n^2 = \infty$$

there exists an orthogonal series

$$(3) \quad \sum_{n=1}^{\infty} a_n \varphi_n(x)$$

which everywhere on  $[0,1]$  is summable with a certain Toeplitz-

method  $T$ , while no subsequence  $S_{k_1}(x) = \sum_{n=1}^{k_1} a_n \varphi_n(x)$  con-

verges in any point  $x \in [0,1]$ .

Theorem 3 : Let  $\{\varphi_n(x)\}$  be a bounded orthogonally normed system on  $[0,1]$ . Then there exists a number  $\delta > 0$  so that

Card 2/4

67508

SOV/155-59-1-11/30

Unconditional Convergence With Respect to  $+\infty$

if the series

$$(22) \quad \sum_{n=1}^{\infty} a_n \varphi_n(x), \quad |\varphi_n(x)| \leq A$$

has partial sums for an arbitrary arrangement of the terms

$$\sum_{i=1}^{\infty} a_{k_i} \varphi_{k_i}(x) \quad \text{satisfying the inequation}$$

$$(23) \quad \lim_{N \rightarrow \infty} \sum_{i=1}^N a_{k_i} \varphi_{k_i}(x) > -\infty \quad \text{for } x \in E,$$

where  $m \in E > 1 - \delta$ , then

$$(24) \quad \sum_{n=1}^{\infty} |a_n| < \infty$$

i.e. the series (22) converges absolutely on  $[0,1]$ . From

X

Card 3/4

67508

15

Unconditional Convergence With Respect to  $+\infty$  SOV/155-59-1-11/30

this theorem there results as a special case a theorem of Privalov [Ref 4].  
Theorem 4: If  $\{\varphi_n(x)\}$  is a bounded orthogonally normed system

on  $[0,1]$ , then there exists no series  $\sum_{n=1}^{\infty} a_n \varphi_n(x)$  which on

a set  $E \subset [0,1]$  with  $m E = 1$  is unconditionally convergent with respect to  $+\infty$ .

Theorem 6': There exists no trigonometric series

$\sum_{n=1}^{\infty} (a_n \cos 2\pi nx + b_n \sin 2\pi nx)$  which on  $E$  with  $m E > 0$  is

unconditionally convergent with respect to  $+\infty$ .

The author mentions Z.N. Kazhdan.

There are 5 references, 3 of which are Soviet, 1 Polish and 1 American.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet imeni M.V. Lomonosova  
(Moscow State University imeni M.V. Lomonosov)

SUBMITTED: January 19, 1959

Card 4/4

UL'YANOV, P.L.

Local properties of convergent Fourier series. Uch.zap.Mosk.  
un. no.186[a]:71-82 '59. (MIRA 13:6)  
(Fourier's series)

69472

16.4200

S/055/59/000/05/004/020

AUTHOR: Ul'yanov, P. L.

TITLE: Singular Integrals and Fourier Series

PERIODICAL: Vestnik Moskovskogo universiteta. Seriya matematiki, mekhaniki, astronomii, fiziki, khimii, 1959, No. 5, pp. 33-42  
vol. 14

TEXT: The author constructs a continuous function  $f(x)$  for which the limit

$$(5) \quad \lim_{h \rightarrow 0} \int_h^\pi \frac{f(x+t) + f(x-t) - 2f(x)}{t} dt$$

exists for no  $x$ . The Fourier series of this function, however, is uniformly convergent. Moreover it is shown that the functions  $f(x)$  with these properties form a set of first category in the set of the continuous  $2\pi$ -periodical functions. Furthermore it is proved: Theorem 2: There exist two conjugate continuous periodical functions  $F_1(x)$  and  $F_2(x)$  with the properties:

$$1.) \quad \int_0^{2\pi} \frac{|F_i(x+t) - F_i(x-t)|}{t} dt = \infty \quad \text{for all } x; i = 1, 2$$

Card 1/2

Singular Integrals and Fourier Series

69472

S/055/59/000/05/004/020

2.)  $\lim_{h \rightarrow 0} \int_h^{\pi} \frac{F_i(x+t) - F_i(x-t)}{t} dt$  exists for all  $x$ ;  $i = 1, 2$

3.) The Fourier series of  $F_1(x)$  and  $F_2(x)$  converge uniformly on  $[0, 2\pi]$ .

The author mentions N. N. Luzin and Kolmogorov.

There are 6 references: 2 Soviet, 3 Polish and 1 English

SUBMITTED: October 12, 1956

Card 2/2

~~16(1)~~ 16.2600

AUTHOR: Ul'yanov, P.L.

05704

SOV/38-23-5-8/8

TITLE: Unconditional Summability

PERIODICAL: Izvestiya Akademii nauk SSSR, Seriya matematicheskaya, 1959,  
Vol 23, Nr 5, pp 781 - 808 (USSR)

ABSTRACT: The paper contains proofs and some generalizations of the questions already treated by the author in [Ref 4,5,6] concerning the unconditional summability of function and numerical series, whereby the notion of summability is somewhat extended. Altogether the author gives eight theorems, eleven conclusions and ten lemmata. He mentions I.I. Volkov and A.M. Olevskiy.

There are 12 references, 6 of which are Soviet, 3 Polish, 2 English, and 1 American.

PRESENTED: by A. N. Kolmogorov, Academician

SUBMITTED: December 7, 1958

Card 1/1



16.4000

35821  
S/044/62/000/002/006/092  
C111/C222

AUTHOR: Ul'yanov, P. L.

TITLE: Convergence and summability

PERIODICAL: Referativnyy zhurnal, Matematika, no. 2, 1962, 12-13,  
abstract 2B59. ("Tr. Mosk. matem. o-va," 1960, 2,  
373-399)

TEXT: This paper is a continuation of the author's examination  
of unconditionally summable (in one sense or another) function series  
(Rzh. Mat., 1960, 7396). By  $B = \| B_{nm} \|$  linear regular summation methods  
with the aid of factors are denoted.  $B^* = \| B_{nm} \|$  denotes methods  
which satisfy the conditions

$$\lim_{n \rightarrow \infty} B_{nm} = 1 \quad (m = 0, 1, 2, \dots), \quad (1).$$

$$\lim_{m \rightarrow \infty} B_{nm} = \gamma_n, \quad \lim_{n \rightarrow \infty} \gamma_n = 0$$

$B^{**}$  denotes methods having matrices which satisfy (1). By  $T^* = \| a_{nm} \|$

Card 1/6

Convergence and summability

S/044/62/000/002/008/092  
C111/C222

linear Toeplitz methods are denoted for which

$$\lim_{n \rightarrow \infty} a_{nm} = 0 \quad (m = 0, 1, \dots), \quad \lim_{n \rightarrow \infty} \sum_{nm} a_{nm} = 1.$$

Function series

$$\sum_{n=0}^{\infty} f_n(x) \quad (x \in E) \quad (2)$$

are considered, where the  $f_n(x)$  may not be measurable. The series

$$\sum_{k=0}^{\infty} f_{n_k}(x)$$

is called a partial series of the first kind of (2), and the series

$$\sum_{n=0}^{\infty} \delta_n f_n(x), \quad \delta_n = 0, \text{ or } 1$$

Card 2/6

*f*

S/044/62/000/002/008/092  
C111/C222

Convergence and summability

is a partial series of the second kind of (2). The series

$$\sum f_{v_k}(x)$$

resulting by rearranging the terms of (2) is called a weak rearrangement of (2) if the sequence  $\{v_k\}$  splits into finitely many increasing sequences. If for every weak rearrangement of (2) the B-means  $G_N(x)$  of the resulting series  $(G_N(x))$  is understood in the sense of convergence with respect to the outer measure) converge for  $N \rightarrow \infty$  on  $E$  (almost everywhere on  $E$ ) with respect to the outer measure, then (2) is weakly, unconditionally B-summable with respect to the outer measure on  $E$  (almost everywhere on  $E$ ). The weak unconditional  $B^{*-}$ ,  $B^{**}$  -, and  $T^{*-}$  summability with respect to the outer measure on  $E$ , or almost everywhere on  $E$ , are defined in analogy.

Theorem 1: If the series

$$\sum_{n=0}^{\infty} \varphi_n(x) \quad (x \in E)$$

is weakly, unconditionally  $B^{**}$  - summable ( $T^{*-}$  summable) on  $E$  with Card  $3/6$

J

Convergence and summability

S/044/62/000/002/008/092  
C111/C222

respect to the outer measure, then

$$\psi_n(x) = f(x) + \eta_n(x), \quad x \in E$$

where  $f(x)$  is a finite function on  $E$ , and the series

$$\sum_{n=0}^{\infty} \eta_n(x)$$

converges unconditionally on  $E$  according to the outer measure. If the method  $B^*$  (method  $T^*$ ) does not sum-up the series

$$\sum_{n=0}^{\infty} 1$$

(3)

then

$$f(x) = 0, \quad x \in E.$$

Theorem 5: If the series

Card 4/6

✓

Convergence and summability

S/044/62/000/002/003/092  
C111/C222

$$\sum_{n=0}^{\infty} \psi_n(x) \quad (x \in [0,1])$$

is such that each of its partial series of the first kind on  $[0,1]$  is  $B^{**}$ -summable with respect to the outer measure, then

$$\psi_n(x) = f(x) + \varphi_n(x)$$

where  $f(x)$  is a finite function, and the series  $\sum_{n=0}^{\infty} \varphi_n(x)$  converges on  $[0,1]$  unconditionally with respect to the outer measure. Here  $f(x) = 0$  if (3) is not  $B^{**}$ -summable.

Theorem 7: If the series

$$\sum_{i=0}^{\infty} f_i(x), \quad x \in E \quad (4)$$

is such that each of its partial series of the second kind is  $B^{**}$ -summable on  $E$  with respect to the outer measure, then (4) is uncondi-  
Card 5/6

f

S/044/62/000/002/008/092  
C111/C222

Convergence and summability

tionally convergent on  $E$  with respect to the outer measure.

A few conclusions are drawn from the stated theorems. The unconditional summability almost everywhere and the case of numerical series are considered. Applications of the obtained results are given regarding orthogonal series and series of the type

$$\sum_{n=0}^{\infty} a_n \varphi(\lambda_n x + \beta_n)$$

where  $\varphi(x)$  is a periodic function, the integral of which is 0.

[Abstracter's note: Complete translation.]

Card 6/6

V

30005  
S/550/60/009/000/004/008  
D251/D305

16.4000

AUTHOR: Ul'yanov, P.L.

TITLE: Convergence and summability

SOURCE: Moskovskoye matematicheskoye obshchestvo Trudy,  
v. 9, 1960, 373 - 399

TEXT: The results of this article were reported to the Moscow  
Mathematical Association on November 24, 1959. The author defines  
 $B = //B_{n,m} //$  as the methods satisfying

$$\lim_{n \rightarrow \infty} B_{n,m} = 1 \quad (m = 0, 1, \dots) \quad (1)$$

and

$$\lim_{n \rightarrow \infty} B_{n,m} = \gamma_n, \quad \lim_{n \rightarrow \infty} \gamma_n = 0. \quad (2)$$

If only (1) is satisfied, the method is denoted by  $B^{**}$ ,  $T^* = a_{n,m}$   
denotes the linear methods of Tepits

Card 1/14

$$\lim_{n \rightarrow \infty} a_{n,m} = 0 \quad (m = 0, 1, \dots) \quad (3)$$

Convergence and summability

30005  
S/550/60/009/000/004/008  
D251/D305

$$\lim_{n \rightarrow \infty} \sum_{m=0}^{\infty} a_{n,m} = 1. \quad (4)$$

The author then states and proves the following theorems: Theorem 1: If the series

$$\sum_{n=0}^{\infty} \psi_n(x) \quad (x \in E) \quad (19)$$

is weakly absolutely B\*\* - summable (T\* summable) on E according to the lower measure that

$$\psi_n(x) = f(x) + \eta_n(x) \quad (x \in E) \quad (20)$$

where f(x) is a finite function on E and

$$\sum_{n=0}^{\infty} \eta_n(x) \quad (21)$$

Card 2/14



30005

S/550/60/009/000/004/008  
D251/D305

# Convergence and summability

is absolutely convergent on E according to the lower measure. Also, if the method B\*\* (T\*) does not sum the series

$$\sum_{n=0}^{\infty} 1 \quad (20)$$

then  $f(x) \equiv 0$  for  $x \in E$ . Theorem 2: If series (19) consists of metric functions and is weakly absolutely B\*\*--summable (T\*-summable) on E according to the measure (20), then series (21) is absolutely convergent on E according to the measure and

$$\sum_{n=0}^{\infty} \eta_n^2(x) < \infty \quad (29)$$

almost everywhere on E. Also, if B\*\* (T\*) does not sum the series (22) then  $f(x) \equiv 0$  on E. Theorem 3: If near the terms of the series

Card 3/14

Convergence and summability

30005  
S/550/60/009/000/004/008  
D251/D305

$$\sum_{n=0}^{\infty} \psi_n(x) \quad (x \in [0, 1]) \quad (30)$$

there are infinitely many metric functions and the series (30) is weakly absolutely B\*-summable (T\*-summable) almost everywhere on  $[0, 1]$ , then

$$\psi_n(x) = f(x) + \eta_n(x), \quad (x \in [0, 1]) \quad (31)$$

where  $f(x)$  is a metric finite function on  $[0, 1]$  and the series

$$\sum_{n=0}^{\infty} \eta_n(x) \quad (32)$$

is weakly absolutely convergent almost everywhere on  $[0, 1]$ . If B\* (T\*) does not sum (22) then  $f(x) \equiv 0$ . The result of A.M. Olevs-kiy (Ref. 15: DAN 125, No. 2, 1959, 269-272) is mentioned in the discussion on this theorem. Theorem 4: There exists a regular me-

Card 4/14

Convergence and summability

30005  
S/550/60/009/000/004/008  
D22/D305

thod  $B = //B_{n,m} //$  and an orthogonal series

$$\sum_{n=0}^{\infty} a_n \varphi_n(x) \quad (a_n \varphi_n(x) \rightarrow 0 \text{ on } [0, 1]) \quad (33)$$

which diverges everywhere on  $[0, 1]$  and which nevertheless is absolutely B-summable almost everywhere on  $[0, 1]$ . The orthogonal series of Men'shov is used in the proof (Ref. 14: Kachmazh S., and G. Shteyngauz, Teoriya ortogonal'nykh ryadov (Theory of Orthogonal Series) M., Fizmatgiz, 1958). Theorem 3: If the series

$$\sum_{n=0}^{\infty} \psi_n(x) \quad (x \in [0, 1]) \quad (48)$$

is such that any of its partial series of the first kind are B\*-summable on  $[0, 1]$  according to the lower measure

$$\psi_n(x) = f(x) + \eta_n(x)$$

Card 5/14

Convergence and summability

30005  
S/550/60/009/000/004/008  
D251/D305

where  $f(x)$  is a finite function and the series

$$\sum_{n=0}^{\infty} \eta_n(x)$$

is absolutely convergent on  $[0, 1]$  according to the lower measure.  $f(x) = 0$  if (22) is not  $B^{**}$ -summable. Theorem 6: There exists an orthogonal series

$$\sum_{n=0}^{\infty} c_n \varphi_n(x) \quad (c_n \varphi_n(x) \rightarrow 0 \text{ on } [0, 1]), \quad (56)$$

and which nevertheless is such that any one of its partial series of the first kind is  $(0, 1)$ -summable almost everywhere on  $[0, 1]$  [Abstractor's note:  $(0, 1)$  summability not defined]. Theorem 7: If the series

Card 6/14

Convergence and summability

30005  
S/550/60/009/000/004/008  
D251/D305

$$\sum_{i=0}^{\infty} f_i(x) \quad (x \in E) \quad (70)$$

is such that any of its partial series of the second kind is  $B^{**}$ -summable on  $E$  according to the lower measure, then (70) is absolutely convergent on  $E$  to the lower measure. Theorem 8: If series (70) consists of metric functions on  $[0, 1]$  and any of its partial series of the second kind is  $B^{**}$ -summable on  $[0, 1]$ , then this series is absolutely convergent on  $[0, 1]$  according to the measure, and

$$\sum_{i=0}^{\infty} f_i^2(x) < \infty \text{ for almost all } x \in [0, 1] \quad (72)$$

Theorem 9: If the series

$$\sum_{i=0}^{\infty} f_i(x) \quad (x \in E) \quad (75)$$

Card 7/14